

12 **EUROPEAN PATENT APPLICATION**

21 Application number: 85306382.4  
 22 Date of filing: 09.09.85


51 Int. Cl.<sup>4</sup>: **C 07 D 405/06**  
**A 61 K 31/40**

30 Priority: 24.09.84 US 653798  
 10.12.84 US 679676  
 43 Date of publication of application:  
 30.04.86 Bulletin 86/18  
 84 Designated Contracting States:  
 AT BE CH DE FR GB IT LI LU NL SE

71 Applicant: WARNER-LAMBERT COMPANY  
 201 Tabor Road  
 Morris Plains New Jersey 07950(US)  
 72 Inventor: Hoeffle, Milton L.  
 1020 Belmont  
 Ann Arbor Michigan 48104(US)  
 72 Inventor: Roth, Bruce D.  
 1440 King George Boulevard  
 Ann Arbor Michigan 48104(US)  
 72 Inventor: Stratton, Charlotte D.  
 1523 Covington Drive  
 Ann Arbor Michigan 48103(US)  
 74 Representative: Jones, Michael Raymond et al,  
 HASELTINE LAKE & CO. Hazlitt House 28 Southampton  
 Buildings Chancery Lane  
 London WC2A 1AT(GB)

84 Trans-6-[2-(substitutedpyrrol-1-yl)alkyl]-pyran-2-one inhibitors of cholesterol synthesis.

87 6-[2-(Substituted-pyrrol-1-yl)alkyl]pyran-2-ones of formula I



and the corresponding ring-opened hydroxy-acids derived therefrom are potent inhibitors of the enzyme 3-hydroxy-3-methylglutarylcoenzyme A reductase (HMG-CoA reductase), and are thus useful hypolipidemic and hypocholesterolemic agents. Pharmaceutical compositions containing such compounds, and a method of preparing the compounds are also disclosed.

EP 0 179 559 A2

-1-

TRANS-6-[2-(SUBSTITUTEDPYRROL-1-YL)ALKYL]-  
PYRAN-2-ONE INHIBITORS OF CHOLESTEROL SYNTHESIS

The present invention is related to compounds and pharmaceutical compositions useful as hypocholesterolemic and hypolipidemic agents. More particularly, this invention concerns certain trans-6-[2-(substitutedpyrrol-1-yl)alkyl]-2-ones and the corresponding ring-opened acids derived therefrom which are potent inhibitors of the enzyme 3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMG-CoA reductase), pharmaceutical composition containing such compounds, and a method of lowering blood serum cholesterol levels employing such pharmaceutical compositions.

15

20

High levels of blood cholesterol and blood lipids are conditions which are involved in the onset of arteriosclerosis. It is well known that inhibitors of HMG-CoA reductase are effective in lowering the level of blood plasma cholesterol, especially low density lipoprotein cholesterol (LDL-C), in man (cf. M. S. Brown and J. L. Goldstein, New England Journal of Medicine (1981), 385, No. 9, 515-517). It has now been established that lowering LDL-C levels affords protection from coronary heart disease (cf. Journal of the American Medical Association (1984) 251, No. 3, 351-374).

Moreover, it is known that certain derivatives of mevalonic acid (3,5-dihydroxy-3-methylpentanoic acid) and the corresponding ring-closed lactone form, mevalonolactone, inhibit the biosynthesis of cholesterol (cf. F. M. Singer et al., Proc. Soc. Exper. Biol. Med. (1959), 102, 278) and F. H. Hulcher, Arch. Biochem. Biophys. (1971), 146, 422).

United States Patents 3,983,148; 4,849,495 and 4,137,322 disclose the fermentative production of a natural product, now called compactin, having an inhibitory effect on cholesterol biosynthesis. Compactin has been shown to have a complex structure which includes a mevalonolactone moiety (Brown et al., J. Chem. Soc. Perkin I, (1976), 1165).

30

35

United States Patent 4,255,444 to Oka et al. 017.9559  
discloses several synthetic derivatives of mevalonolactone  
having antilipidemic activity.


United States Patents 4,198,425 and 4,262,813 to  
5 Mitsue et al. disclose aralkyl derivatives of mevalono-  
lactone which are useful in the treatment of hyperlipid-  
emia.

United States Patent 4,375,475 to Willard et al.  
discloses certain substituted 4-hydroxytetrahydropyran-  
10 2-ones which, in the 4(R)-trans stereoisomeric form, are  
inhibitors of cholesterol biosynthesis.

15 In accordance with the present invention, there are  
provided certain trans-6-[2-(substitutedpyrrol-1-yl)-  
alkyl]pyran-2-ones and the corresponding ring-opened  
hydroxy-acids derived therefrom which are potent inhibi-  
tors of cholesterol biosynthesis by virtue of their  
20 ability to inhibit the enzyme 3-hydroxy-3-methylglutaryl-  
coenzyme A reductase (HMG-CoA reductase).

In particular, in its broadest chemical compound  
aspect, the present invention provides compounds of  
structural formula I


25



30


wherein X is -CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>-, or -CH(CH<sub>3</sub>)CH<sub>2</sub>-. R<sub>1</sub> is  
1-naphthyl; 2-naphthyl; cyclohexyl; norbornenyl;  
35 phenyl; phenyl substituted by fluorine, chlorine,  
hydroxy, trifluoromethyl, alkyl of from one to four  
carbon atoms, alkoxy of from one to four carbon

atoms, or alkanoyloxy of from <sup>-4-</sup>two to eight carbon atoms;  
 2-, 3-, or 4-pyridinyl; 2-, 3-, or 4-pyridinyl-N-oxide;  
 or




where  $R_5$  is alkyl of from one to four carbon atoms and  
 $\text{hal}^-$  is chloride, bromide, or iodide.  $R_2$  and  $R_3$  are  
 independently hydrogen; chlorine; bromine; cyano;  
 10 trifluoromethyl; phenyl; alkyl of from one to four carbon  
 atoms; carboalkoxy of from two to eight carbon atoms;  
 $-\text{CH}_2\text{OR}_6$  where  $R_6$  is hydrogen, alkanoyl of from one to six  
 carbon atoms, or where  $R_2$  and  $R_3$  are  $-\text{CH}_2\text{OCONHR}_7$  where  $R_7$   
 is alkyl of from one to six carbon atoms, phenyl, or  
 15 phenyl substituted with chlorine, bromine, or alkyl of  
 from one to four carbon atoms.  $R_2$  and  $R_3$  may also, when  
 taken together with the carbon atoms to which they are  
 attached, form a ring denoted by

20




where  $n$  is three or four; a ring denoted by

25




a ring denoted by

30



35


where  $R_8$  is hydrogen, alkyl of from one to six carbon atoms, phenyl, or benzyl; or a ring denoted by



where  $R_9$  and  $R_{10}$  are hydrogen, alkyl of from one to four carbon atoms, or benzyl.

$R_4$  is alkyl of from one to four carbon atoms, cyclopropyl, cyclobutyl, or trifluoromethyl.


Also contemplated as falling within this aspect of the invention are the corresponding dihydroxy-acid compounds of formula II corresponding to the opened form of the lactone ring of compounds of formula I



II


where  $X$ ,  $R_1$ ,  $R_2$ ,  $R_3$ , and  $R_4$  are as defined above, and the pharmaceutically acceptable salts thereof, all of the compounds being in the trans racemate of the tetrahydropyran moiety.

In another aspect of the present invention, there is provided a method of preparing compounds of formula I above by (a) first reacting a substituted [(pyrrol-1-yl)-alkyl]aldehyde compound of formula III



III

where  $X$ ,  $R_1$ ,  $R_2$ ,  $R_3$ , and  $R_4$  are as defined above, with the alkali metal salt of the dianion of methyl acetoacetate to form a compound of structural formula IV




0179559

5

IV

where X, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are as defined above, then successivly (b) reducing compound IV with a trialkylborane and sodium borohydride and (c) oxidizing with alkaline hydrogen peroxide to produce an acid compound of formula V

15



V

and finally (d) cyclizing, if desired, the acid compound of formula V to a lactone compound of formula I by heating in an inert solvent or, alternatively converting, if desired, the acid compound of formula V to a pharmaceutically acceptable salt.

In another aspect, the present invention provides pharmaceutical compositions, useful as hypolipidemic or hypocholesterolemic agents, comprising a hypolipidemic or hypocholesterolemic affective amount of a compound in accordance with this invention as set forth above, in combination with a pharmaceutically acceptable carrier.


In another aspect, the present invention provides a method of inhibiting cholesterol biosynthesis in a patient in need of such treatment by administering a pharmaceutical composition in accordance with the present invention as defined above.

35

In a first preferred subgeneric chemical compound aspect, the present invention provides compounds of formula I above wherein X is -CH<sub>2</sub>CH<sub>2</sub>-, R<sub>1</sub> is

as defined above,  $R_2$  and  $R_3$  are independently hydrogen, chlorine, or bromine, and  $R_4$  is as defined above.

In a second preferred subgeneric chemical compound aspect, the present invention provides compounds of formula I above where X is  $-\text{CH}_2\text{CH}_2-$ ,  $R_1$  is  
 5 phenyl or phenyl substituted by fluorine, chlorine, hydroxy, trifluoromethyl, alkyl of from one to four carbon atoms, alkoxy of from one to four carbon atoms, or alkanoyloxy of from two to eight carbon atoms, or where  $R_1$  is 2-, 3-, or 4-pyridinyl; 2-, 3-, or  
 10 4-pyridinyl-N-oxide, or




15

where  $R_5$  is alkyl of from one to four carbon atoms and  $\text{hal}^-$  is chloride, bromide, or iodide. In this aspect of the invention,  $R_2$  and  $R_3$  are preferably independently hydrogen, chlorine, or bromine, and  $R_4$  is alkyl of from  
 20 one to four carbon atoms or trifluoromethyl.


In a third preferred subgeneric chemical compound aspect, the present invention provides compounds of formula I above where X is  $-\text{CH}_2\text{CH}_2-$ ,  $R_1$  is phenyl or phenyl substituted by fluorine, chlorine, hydroxy,  
 25 trifluoromethyl, alkoxy of from one to four carbon atoms, or alkanoyloxy of from two to eight carbon atoms,  $R_2$  and  $R_3$  are independently hydrogen, chlorine, or bromine, and  $R_4$  is isopropyl or trifluoromethyl.

30 In a fourth preferred subgeneric chemical compound aspect, the present invention provides compounds of formula I above where X is  $-\text{CH}_2\text{CH}_2-$ , and  $R_1$  is phenyl or phenyl substituted by fluorine, chlorine, trifluoromethyl, alkyl of from one to four  
 35 carbon atoms, alkoxy of from one to four carbon atoms, or alkanoyloxy of from two to eight carbon atoms, or where  $R_1$  is 1-naphthyl, or 2-naphthyl. In this preferred aspect of the invention,  $R_2$  and  $R_3$  are independently


hydrogen, chlorine, bromine, cyano, trifluoromethyl, phenyl, alkyl of from one to four carbon atoms, carboalkoxy of from two to eight carbon atoms,  $-\text{CH}_2\text{OR}_6$  where  $\text{R}_6$  is hydrogen or alkanoyl of from one to six carbon atoms,  $-\text{CH}_2\text{OCONHR}_7$  where  $\text{R}_7$  is alkyl of from one to six carbon atoms, phenyl, or phenyl substituted with chlorine, bromine, or alkyl of from one to four carbon atoms. In this aspect of the invention,  $\text{R}_2$  and  $\text{R}_3$  may also, when taken together with the carbon atoms to which they are attached, form a ring denoted by




where  $n$  is three or four; a ring denoted by



a ring denoted by



where  $\text{R}_8$  is hydrogen, alkyl of from one to four carbon atoms, phenyl, or benzyl; or a ring denoted by




where  $\text{R}_9$  and  $\text{R}_{10}$  are hydrogen, alkyl of from one to four carbon atoms, or benzyl. In this aspect of the invention,  $\text{R}_4$  is preferably alkyl of from one to four carbon atoms, cyclopropyl, cyclobutyl, or trifluoro-

methyl.


In a fifth preferred subgeneric chemical compound aspect, the present invention provides compounds of formula I above where  $\text{X}$  is  $-\text{CH}_2\text{CH}_2-$ , and  $\text{R}_1$

is phenyl or phenyl substituted by fluorine, chlorine, trifluoromethyl, alkyl of from one to four carbon atoms, alkoxy of from one to four carbon atoms, or alkanoyloxy of from two to eight carbon atoms.  $R_2$  and  $R_3$  are preferably independently hydrogen, chlorine, bromine, phenyl, or carboalkoxy of from two to eight carbon atoms. In this aspect of the invention  $R_2$  and  $R_3$  may also, when taken together with the carbon atoms to which they are attached, form a ring denoted by


18



15 where  $n$  is three or four; a ring denoted by



20 where  $R_8$  is hydrogen, or alkyl of from one to four carbon atoms; or a ring denoted by




25

where  $R_9$  and  $R_{10}$  are hydrogen or alkyl of from one to four carbon atoms. In this aspect of the invention,  $R_4$  is preferably alkyl of from one to four carbon atoms, or trifluoromethyl.

In a sixth preferred subgeneric chemical compound aspect, the present invention provides compounds of formula I above where  $X$  is  $-\text{CH}_2\text{CH}_2-$ ,  $R_1$  is phenyl or phenyl substituted by fluorine, chlorine, trifluoromethyl, alkyl of from one to four carbon atoms, alkoxy of from one to four carbon atoms, or alkanoyloxy of from two to eight carbon atoms.  $R_2$  and  $R_3$  are

preferably independently carboalkoxy of from two to eight carbon atoms or, when taken together with the carbon atoms to which they are attached form a ring denoted by

5



10 where R<sub>8</sub> is hydrogen or alkyl of from one to four carbon atoms. In this aspect of the invention, R<sub>4</sub> is preferably isopropyl or trifluoromethyl.

As used throughout this specification and the appended claims, the term "alkyl" denotes a branched or  
15 unbranched saturated hydrocarbon group derived by the removal of one hydrogen atom from an alkane.

The term "alkoxy" denotes an alkyl group, as just defined, attached to the parent molecular residue through an oxygen atom.

20 The term "alkanoyloxy" is meant to denote an alkyl group, as defined above, attached to a carbonyl group and thence, through an oxygen atom, to the parent molecular residue.

The term "carboalkoxy" is meant to denote an alkyl  
25 group, as defined above, attached to an oxygen atom and thence, through a carbonyl group, to the parent molecular residue.

The term "norbornenyl" denotes a group derived by the removal of a hydrogen atom (other than at a  
30 bridgehead carbon atom) from bicyclo[2.2.1]hept-2-ene.

---

- // -

Specific examples of compounds contemplated as falling within the scope of the present invention include the following:

- 5     trans-6-[2-[2-Cyclobutyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.
- trans-6-[2-[2-Cyclohexyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-pyran-2-one.
- trans-Tetrahydro-4-hydroxy-6-[2-(2-methyl-5-phenyl-1H-pyrrol-1-yl)ethyl]-2H-pyran-2-one.
- 10    trans-6-[2-[2-(4-Chlorophenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.
- trans-Tetrahydro-4-hydroxy-6-[2-[2-(4-methoxyphenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one.
- trans-6-[2-[2-([1,1'-Biphenyl]-4-yl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.
- 15    trans-Tetrahydro-4-hydroxy-6-[2-[2-methyl-5-[3-(trifluoromethyl)phenyl]-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one.
- trans-6-[2-[2-(2,5-Dimethylphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.
- 20    trans-6-[2-[2-(2,6-Dimethoxyphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.
- 25    trans-Tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(2-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one.
- trans-6-[2-(2-(Cyclohexyl-5-trifluoromethyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.
- trans-6-[2-[2-(4-Fluorophenyl)-3,4-dimethyl-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.
- 30    trans-2-(4-Fluorophenyl)-5-(1-methylethyl)-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3,5-dicarboxylic acid.
- 35    trans-2-(4-Fluorophenyl)-N<sup>3</sup>,N<sup>3</sup>,N<sup>4</sup>,N<sup>4</sup>-tetramethyl-5-(1-methylethyl)-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3,4-dicarboxamide.

trans-6-[2-[3,4-Dichloro-2-(3-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

5       trans-2-(4-Fluorophenyl)-5-(1-methylethyl)-1-[2-(tetrahydro)-4-hydroxy-6-oxo-2H-pyran-2-yl]ethyl]-1H-pyrrole-3,4-dicarbonitrile.

trans-6-[2-[3,4-Diacetyl-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

10       trans-Diethyl 2-(4-Fluorophenyl)-1-[2-(tetrahydro)-4-hydroxy-6-oxo-2H-pyran-2-yl]ethyl]-5-(trifluoromethyl)-1H-pyrrole-3,4-dicarboxylate.

15       trans-Bis(1-methylethyl) 2-(4-Fluorophenyl)-5-(1-methylethyl)-1-[2-(tetrahydro)-4-hydroxy-6-oxo-2H-pyran-2-yl]ethyl]-1H-pyrrole-3,4-dicarboxylate.

trans-6-[2-[3,4-Diethyl-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

20       trans-6-[2-[2-(4-Fluorophenyl)-3,4-bis(hydroxymethyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-1-Methylethyl 4-Chloro-2-(4-fluorophenyl)-5-(1-methylethyl)-1-[2-(tetrahydro)-4-hydroxy-6-oxo-2H-pyran-2-yl]ethyl]-1H-pyrrole-3-carboxylate.

25       trans-6-[2-[4-(4-Fluorophenyl)-6-(1-methylethyl)-1H-furo[3,4-c]pyrrol-5(3H)-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

30       trans-6-[2-[2-(4-Fluorophenyl)-5-(1-methylethyl)-3,4-bis[[[(phenylamino)carbonyl]oxy]methyl]-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-1-Methylethyl 4-Chloro-5-(4-fluorophenyl)-2-(1-methylethyl)-1-[2-(tetrahydro)-4-hydroxy-6-oxo-2H-pyran-2-yl]ethyl]-1H-pyrrole-3-carboxylate.

35       trans-Ethyl 5-(4-Fluorophenyl)-1-[2-(tetrahydro)-4-hydroxy-6-oxo-2H-pyran-2-yl]ethyl]-2-(trifluoromethyl)-1H-pyrrole-3-carboxylate.

trans-Ethyl 5-(4-Fluorophenyl)-2-(1-methylethyl)-4-phenyl-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxylate.

5 trans-6-[2-[1-(4-Fluorophenyl)-4,5,6,7-tetrahydro-3-methyl-2H-isoindol-2-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-4-(4-Fluorophenyl)-2-methyl-6-(1-methylethyl)-5-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-pyrrolo[3,4-c]pyrrole-1,3(2H,5H)-dione.

10 trans-6-[2-[1-(4-Fluorophenyl)-5,6-dihydro-3-(1-methylethyl)pyrrolo[3,4-c]pyrrol-2(4H)-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

15 trans-6-[2-[1-(4-Fluorophenyl)-5,6-dihydro-5-methyl-3-(1-methylethyl)pyrrolo[3,4-c]pyrrol-2(4H)-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-6-[2-[3-Chloro-5-(4-fluorophenyl)-2-(1-methylethyl)-4-phenyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

20 trans-6-[2-[2-(4-Fluorophenyl)-5-(1-methylethyl)-3,4-diphenyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

Particularly preferred compounds in accordance with the present invention are:

25 trans-6-[2-[3,4-Dichloro-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-6-[2-[3,4-Dibromo-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

30 trans-6-[2-[2-(4-Fluorophenyl)-5-(trifluoromethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-Dimethyl 2-(4-Fluorophenyl)-5-(1-methylethyl)-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3,4-dicarboxylate.

35 trans-6-[2-[2-(4-Fluorophenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-6-[2-[2-(4-Fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-6-[2-[2-Cyclopropyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

5     trans-6-[2-[2-(1,1-Dimethylethyl)-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-Tetrahydro-4-hydroxy-5-[2-[2-(2-methoxyphenyl)-5-trifluoromethyl-1H-pyrrol-1-yl]ethyl]-2H-2-one.

10     trans-Tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one.

trans-Tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(1-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one.

15     trans-6-[2-(2-Bicyclo[2.2.1]hept-5-en-2-yl-5-methyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

trans-6-[2-[2-(4-Fluorophenyl)-5-(1-methylphenyl)-1H-pyrrol-1-yl]propyl]tetrahydro-4-hydroxy-2H-pyran-2-one.





20     Compounds of the present invention where  $R_2$  and  $R_3$  are hydrogen are prepared by the methods outlined in Reaction Sequence 1 or Reaction Sequence 2.  
As shown in Reaction Sequence 1, the aldehydes, VI, are reacted with the appropriately substituted vinylketones,  
25     VII, in the presence of the thiazolium salt, VIII, and a base such as triethylamine, to produce the diketones, IX.  
(See Ang. Chem. Int. Ed., 15: 639-712 (1976)).

The diketones, IX, are reacted with an omega-amino-alkylnitrile (compound Roman numeral ten where the value  
30     of X is methylene, ethylene, or 1-methylethylene) in acetic acid to produce the disubstituted pyrrole nitriles, XI.




Treatment of the pyrrole nitriles, XI, with diisobutylaluminum hydride in an inert solvent such as  
35     dichloromethane produces the corresponding pyrrole aldehydes, XII.

REACTION SEQUENCE 1



0179559



Diisobutylaluminum  
hydride



- 1) Tributylborane
- 2) Sodium borohydride
- 3)  $\text{H}_2\text{O}_2$ ,  $\text{OH}^-$




I

Reaction of the pyrrole aldehydes, XII, ~~Q1h78559~~  
dilithium or lithium sodium salt methyl acetoacetate  
produces the 7-(substitutedpyrrolyl)-5-hydroxy-3-oxo-  
heptanoates, XIII. The heptanoates, XIII, are dissolved  
5 in a polar solvent such as tetrahydrofuran, through which  
a small quantity of air has been bubbled. A slight  
excess of a trialkylborane, such as tributylborane, is  
added to the mixture which is then cooled to a  
temperature of preferably between about 0°C and -78°C  
10 after which sodium borohydride is added.

After stirring this mixture for about one to two  
hours, the mixture is oxidized with basic hydrogen  
peroxide. The reaction produces the 7-(substituted-  
pyrrolyl)-3,5-dihydroxyheptanoic acids, XIV, in which the  
15 product contains a predominance of the desired R\*, R\*  
configuration at carbon atoms three and five which bear  
the hydroxy groups.


The acids may be converted to a corresponding  
pharmaceutically acceptable salt by conventional methods  
20 or, alternatively, cyclized to the 6-[2-(substituted-  
pyrrol-1-yl)alkyl]pyran-2-ones, I, by dehydration in an  
inert solvent such as refluxing toluene with azeotropic  
removal of water. This cyclization reaction is found to  
produce material containing from 85-98% of the desired  
25 active trans-configuration of the 4-hydroxy group  
relative to the 6-(substitutedpyrrolyl)alkyl group on the  
pyran-2-one lactone ring.

Alternative procedures for preparing compounds of  
formula I of this invention where R<sub>2</sub> and R<sub>3</sub> are hydrogen,  
30 and for preparing intermediates, are illustrated in  
Reaction Sequence 2. As shown in Reaction Sequence 2,  
the diketones, IX, can be prepared by reacting the known  
alpha-haloketones, XV, with the sodium salt of known  
beta-ketoesters, XVI, followed by hydrolysis and  
35 decarboxylation in the conventional manner. The  
diketones, IX, are reacted with ammonium acetate in  
acetic acid to produce the cyclized 2,5-disubstituted  
pyrroles, XVII.



XV


XVI



XVIII

$\text{NH}_4\text{OAc}$

$\text{HOAc}$




IX

1)  $\text{NaH}$


2)  $\text{Br-X-CH}(\text{OCH}_3)_2$  (XXI)

3)  $\text{H}^+$ ,  $\text{H}_2\text{O}$

$\text{H}_2\text{N-X-OH}$  (XVIII)



(XXII in  
Reaction  
Sequence 1)




XX (XIII in Reaction  
Sequence 1)

1) Tosyl chloride

2)  $\text{CN}^-$

IX

$\text{H}_2\text{N-X-OH}$   
XVIII



XIX

An alternative for this step, preferred when  $R_1$  and/or  $R_4$  are sterically bulky groups, involves reaction of the diketones, IX, with an omega-hydroxyalkyl amine (compound XVIII where X is methylene, ethylene, 1-methylethylene), to produce the N-(omega-hydroxy-alkyl)-2,5-disubstitutedpyrroles, XIX.

The 2,5-disubstitutedpyrroles, XVII, are converted to the omega-(substitutedpyrrolyl)aldehydes, XX, by sequential reaction with sodium hydride, a 1,1-dimethoxy-omega-bromoalkane (compound XXI where X is methylene, ethylene, 1-methylethylene, or vinyl), and then acid. The aldehydes, XX, are subsequently used in the preparation of compounds of formula I of this invention as illustrated above in Reaction Sequence 1.



The 2,5-disubstituted pyrroles, XVII, are converted to the corresponding (2,5-disubstitutedpyrrolyl)nitriles, XXII (when X is ethylene), by reaction with acrylonitrile or, alternatively (when X is other than ethylene), by starting with compounds of formula XIX. In this latter instance, the hydroxy functionality of compounds of formula XIX is converted to the p-toluenesulfonate by conventional means, and the tosylate group is subsequently displaced by cyanide ion to produce the nitriles of formula XXII. The compounds of formula XXII are subsequently used in the preparation of compounds of formula I of this invention by methods detailed in Reaction Sequence 1 above.

Starting materials and intermediates employed in Reaction Sequences 1' and 2 above may be prepared by the general methods outlined in Reaction Sequence 3. For example, as shown there, the vinyl ketones, VII, are prepared by either of the two methods illustrated. In one method, the known acid chlorides, XXIII, are reacted with the trimethylsilylethene, XXIV, in the presence of anhydrous aluminum chloride in dichloromethane.

In the alternative method of preparing the vinyl ketones, VII, which is preferred when  $R_1$  is an aromatic substituent such as phenyl or substituted phenyl, the

REACTION SEQUENCE 3

0179559



known methyl aryl ketones, XXV, are converted to (dimethylaminoethyl)aryl ketones, XXVI, and then by deamination to the vinyl ketones, VII.


5 The compounds of the present invention of formula I where the groups  $R_2$  and  $R_3$  are other than hydrogen or halogen can be synthesized by the methods detailed in Reaction Sequences 4-8.

Employing the method detailed in Reaction Sequence 4 the compounds of the present invention where  $R_2$  and  $R_3$  are both halogen can be prepared by the halogenation of the unsubstituted compounds with N-halosuccinimide in a three-step process involving the prior protection of the 4-hydroxy group of the lactone ring. Thus, for example, the 2,5-disubstitutedpyrrol-1-yl compounds, XXVII, are first converted to the corresponding tert-butyl-dimethylsilyl ethers, XXVIII. The protected compounds and then chlorinated with N-chlorosuccinimide in a polar solvent such as dimethylformamide to produce the silylated 3,4-dichloro compounds, XXIX. The protecting silyl ether group is then subsequently removed by reaction with a buffered fluoride reagent such as tetrabutylammonium fluoride in a mixed acetic acid/tetrahydrofuran solvent system to produce the dichloro compounds, XXX.


25 Alternatively, as detailed in Reaction Sequence 5, the (2,5-disubstitutedpyrrol-1-yl)alkyl nitriles, XI (see Reaction Sequence 1) are halogenated by employing an N-halosuccinimide in dimethylformamide to provide the 2,5-disubstituted-3,4-dihalopyrroles, XXXI. (See Aiello, et al., J. Het. Chem., 19: 977 (1982)). These compounds can then be subsequently converted to the compounds of the present invention by conventional methods detailed in Reaction Sequence 1.

35 A third method takes advantage of the chemistry of mesionic compounds of the type described originally by R. Huisgen, et al., Ang. Chem. Int. Ed., 3: 136 (1964). In this procedure, detailed in Reaction Sequence 6, an

REACTION SEQUENCE 4 0179559



REACTION SEQUENCE 5




N-alkyl-N-acylamino acid is treated with an acid anhydride and a substituted acetylenic compound to produce a pyrrole. For example, Reaction Sequence 6 shows how reaction of an alpha-halo ester, XXXII, with  
5 2-(1-(2-aminoethyl))-1,3-dioxalane in triethylamine provides the N-alkyl-alpha-aminoester, XXXIII. The aminoester, XXXIII is acylated with an acid chloride and subsequently hydrolyzed in base to produce the N-acyl-N-alkyl aminoacid, XXXIV. Reaction of this latter  
10 compound with the desired substituted acetylenic compound, XXXV, produces the substituted pyrrole compounds, XXXVI. Acidic hydrolysis of XXXVI yields the aldehyde compounds, XXXVII, analogous to compounds XII of Reaction Sequence 1. Compounds of formula XXXVII are  
15 used in subsequent steps in a manner detailed in Reaction Sequence 1 to produce compounds of the present invention.


Preferred substituents for the substituted acetylenic compounds in this method of making compounds of the present invention include carboalkoxy groups,  
20 phenyl groups, alkanoyl groups, alkyl groups and cyano groups. The reaction between the disubstituted acetylene compound and the N-acyl-N-alkyl aminoacids, XXXIV, generally proceeds smoothly; for example, the  
25 dicarbomethoxy acetylene reacts smoothly at 25°C. However, when only one activating group is attached to the acetylene, the reaction mixture must generally be warmed to 70-110°C to obtain high yields of the pyrrole compounds.

30 A variety of other pyrroles can be derived from compounds of the general formula XXXVI when the groups  $R_2$  and  $R_3$  are carbomethoxy. Some of these transformations are detailed in Reaction Sequences 7 and 8. For example, as shown in Reaction Sequence 7, reduction of XXXVI with  
35 a reducing agent such as lithium aluminum hydride results in the bis(hydroxymethyl)pyrrole which can be subsequently further reduced to the dimethyl compound,

REACTION SEQUENCE 6



REACTION SEQUENCE 7



XXXVIII, by means of triethylsilane and trifluoroacetic acid employing the procedure of West, et al., J. Org. Chem., 38: 2675 (1973)).

Alternatively, as shown in Reaction Sequence 8,  
5 reaction of the compounds of formula XXXVI with a Grignard reagent or an alkyl-lithium reagent in the conventional manner followed by reduction and standard work-up affords the higher dialkylpyrroles, XXXIX.

10 Reaction of the diesters, XXXVI, or the corresponding diacids (obtained by conventional hydrolysis) with secondary amines provides the bis(dialkylamides), XL.



Alternatively, reaction of XXXVI with primary  
15 amines, followed by thermal cyclization in the conventional manner, provides the pyrrolosuccinimides, XLI, which can be reduced to XLII, if desired by reducing agents such as lithium aluminum hydride.

The bis(hydroxymethyl)pyrrole compounds derived from  
20 the lithium aluminum hydride reduction of XXXVI can be converted to their corresponding esters or carbamates by reaction with the desired acid anhydride or isocyanate, respectively. (See Anderson, et al., J. Med. Chem., 22: 977 (1979)).

25 The acids, XLIII, derived by convention hydrolysis of compounds of formula XXXVI can also be converted to the bis(amido)pyrroles, XLIV, which in turn can be dehydrated to produce the bis(nitrilo)pyrroles, XLV. Lastly, if desired, the bis(alkanoyl)pyrroles, XLVI, can  
30 be derived from the bis(nitrilo)pyrroles by reaction in the convention manner with the appropriate Grignard reagents.

The ring-opened dihydroxy-acids of structural  
formula II above are intermediates in the synthesis of  
35 the lactone compounds in accordance with the above-detailed reaction methods, or may be produced from the lactone compounds by conventional hydrolysis of the lactone compounds of formula I.

(Throughout this sequence,



In the ring-opened dihydroxy acid form, compounds of the present invention react to form salts with pharmaceutically acceptable metal and amine cations formed from organic and inorganic bases.

5       The term "pharmaceutically acceptable metal cation" contemplates positively charged metal ions derived from sodium, potassium, calcium, magnesium, aluminum, iron, zinc and the like.

10       The term "pharmaceutically acceptable amine cation" contemplates the positively charged ions derived from ammonia and organic nitrogenous bases strong enough to form such cations. Bases useful for the formation of pharmaceutically acceptable nontoxic base addition salts of compounds of the present invention form a class whose  
15 limits are readily understood by those skilled in the art.

20       The free acid form of the compound may be regenerated from the salt, if desired, by contacting the salt with a dilute aqueous solution of an acid such as hydrochloric acid.

25       The base addition salts may differ from the free acid form of compounds of this invention in such physical characteristics as melting point and solubility in polar solvents, but are considered equivalent to the free acid forms for purposes of this invention.

30       The compounds of this invention can exist in unsolvated as well as solvated forms. In general, the solvated forms, with pharmaceutically acceptable solvents such as water, ethanol, and the like, are equivalent to the unsolvated forms for purposes of this invention.

35       The compounds of this invention are useful as hypocholesterolemic or hypolipidemic agents by virtue of their ability to inhibit the biosynthesis of cholesterol through inhibition of the enzyme 3-hydroxy-3-methyl-glutaryl-coenzyme A reductase (HMG-CoA reductase).

      The ability of compounds of the present invention to inhibit the biosynthesis of cholesterol was measured by

two methods. A first method (designated CSI screen) utilized the procedure described by R. E. Dugan et al., Archiv. Biochem. Biophys., (1972), 152, 21-27. In this method, the level of HMG-CoA enzyme activity in standard  
5 laboratory rats is increased by feeding the rats a chow diet containing 5% cholestyramine for four days, after which the rats are sacrificed.

The rat livers are homogenized, and the incorporation of cholesterol-<sup>14</sup>C-acetate into non-  
10 saponifiable lipid by the rat liver homogenate is measured. The micromolar concentration of compound required for 50% inhibition of sterol synthesis over a one-hour period is measured, and expressed as an IC<sub>50</sub> value.


15 A second method (designated COR screen) employed the procedure detailed by T. Kita, et al., J. Clin. Invest., (1980), 66: 1094-1100. In this method, the amount of <sup>14</sup>C-HMG-CoA converted to <sup>14</sup>C-mevalonate in the presence of a purified enzyme preparation of HMG-CoA reductase was  
20 measured. The micromolar concentration of compound required for 50% inhibition of cholesterol synthesis was measured and recorded as an IC<sub>50</sub> value.

The activity of several representative examples of compounds in accordance with the present invention  
25 appears in Table 1, and is compared with that of the prior art compound, compactin. In particular, compounds of the present invention where R<sub>2</sub> and R<sub>3</sub> are substituents other than hydrogen have activities comparable to that of the natural product, compactin.

30

35

TABLE 1



Compound	X	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	IC <sub>50</sub> (Micromoles/Liter) CSI COR
1	-CH <sub>2</sub> CH <sub>2</sub> -	4-Fluorophenyl	H	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	0.48 0.28
2	-CH <sub>2</sub> CH <sub>2</sub> -	4-Fluorophenyl	Cl	Cl	-CH(CH <sub>3</sub> ) <sub>2</sub>	0.16 0.024
3	-CH <sub>2</sub> CH <sub>2</sub> -	4-Fluorophenyl	Br	Br	-CH(CH <sub>3</sub> ) <sub>2</sub>	0.22 0.001
4	-CH <sub>2</sub> CH <sub>2</sub> -	4-Fluorophenyl	-COOCH <sub>3</sub>	-COOCH <sub>3</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	0.11 0.000
5	Compactin (prior art)					0.026 0.028

\* Adjusted for a standard IC<sub>50</sub> value for compactin which was used as an internal standard in the test.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include  
5 powders, tablets, dispersable granules, capsules, cachets, and suppositories.

A solid carrier can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders,  
10 or tablet disintegrating agents; it can also be an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with finely divided active compound. In tablets, the active compound  
15 is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

For preparing suppository preparations, a low-melting wax such as a mixture of fatty-acid glycerides  
20 and cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds and allowed to cool and solidify.

25 The powders and tablets preferably contain 5 to about 70% of the active ingredient. Suitable solid carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, tragacanth, methyl cellulose, sodium  
30 carboxymethyl cellulose, a low-melting wax, cocoa butter, and the like.

The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier providing a  
35 capsule in which the active component (with or without

other carriers) is surrounded by a carrier, which is thus in association with it. Similarly, cachets are included. Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

Liquid form preparations include solutions, suspensions, and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection. Liquid preparations can also be formulated in solution in aqueous polyethylene glycol solution. Aqueous solutions for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavoring agents, stabilizers, and thickening agents as desired. Aqueous suspensions for oral use can be made by dispersing the finely divided active component in water with viscous material, i.e., natural or synthetic gums, resins, methyl, cellulose, sodium carboxymethyl cellulose, and other well-known suspending agents.

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, for example, packeted tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet, or tablet itself or it can be the appropriate number of any of these packaged forms.

In therapeutic use as hypolipidemic or hypocholesterolemic agents, the compounds utilized in the pharmaceutical method of this invention are

administered to the patient at dosage levels of from  
40 mg to 600 mg per day. For a normal human adult  
of approximately 70 kg or body weight, this trans-  
lates to a dosage of from about 0.5 mg/kg to about  
5 8.0 mg/kg of body weight per day.

The dosages, however, may be varied depending  
upon the requirements of the patient, the severity  
of the condition being treated, and the compound  
being employed. Determination of optimum dosages  
10 for a particular situation is within the skill of  
the art.

The following examples illustrate particular  
methods for preparing compounds in accordance with  
this invention. These examples are illustrative  
15 and are not to be read as limiting the scope of the  
invention as it is defined by the appended claims.

#### EXAMPLE 1

Preparation of trans-6-[2-[2-(4-fluorophenyl)-5-  
(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-  
20 4-hydroxy-2H-pyran-2-one

Step A: Preparation of 1-(4-fluorophenyl)-5-  
methyl-1,4-hexanedione.

A mixture of 1-(4-fluorophenyl)-2-propene-1-  
one (43 g, 286.7 mmol) prepared in accordance with  
25 the method detailed in Org. Syn., Coll. Vol. IV,  
305, was mixed with 31.2 ml (344 mmol) of iso-  
butraldehyde, 28 ml (200 mmol) of triethylamine,  
and 14.5 g (57.7 mmol) of 2-(2-hydroxyethyl)-3-  
methyl-4-benzylthiazolium chloride and the mixture  
30 stirred under nitrogen at 70°C for 12 hours.

---

-32-

After this time, the mixture was cooled to room temperature and the cooled mixture was partitioned between ether (500 ml) and water (100 ml). The aqueous layer was further extracted with 300 ml of  
5 ether, the ether solutions combined and washed successively with 200 ml of water, two 200-ml portions of 2M hydrochloric acid, and 100 ml of brine, and finally dried over anhydrous magnesium sulfate.

10 The ether was removed, and the residue was distilled (bp 115-120°C, 0.2 mm Hg) to provide 36.7 g (165 mmol, 58% of 1-(4-fluorophenyl)-5-methyl-1,4-hexanedione which solidified upon standing.

15 Alternate Step A: Preparation of 1-(4-fluorophenyl)-5-methyl-1,4-hexanedione.

Isopropyl vinyl ketone (1.97 g, 20 mmol), prepared from isobutyryl chloride and vinyl trimethylsilane in accordance with the method detailed in  
20 Tet. Letters, (1979), 1995, was mixed with 4-fluorobenzaldehyde (2.4 g, 20 mmol), 2 ml (14 mmol) of triethylamine, and 1.0 g (4.0 mmol) of 2-(2-hydroxyethyl)-3-methyl-4-benzylthiazolium chloride. The mixture was stirred and heated under nitrogen for  
25 five hours. After cooling to room temperature, the mixture was partitioned between ether (200 ml) and water (50 ml). The water layer was extracted with 200 ml of ether and the ether solutions were combined. The combined ether solution was washed  
30 successively with 50 ml of water, two 50-ml portions of 2M hydrochloric acid, and 50 ml of brine. The ether solution was dried over anhydrous magnesium sulfate. After removal of the ether, the remaining liquid was flash chromatographed on silica gel

eluting with 20:1 (volume/volume) hexane-ethyl acetate.  
This procedure afforded 2.59 g of pure 1-(4-fluorophenyl)-5-methyl-1,4-hexanedione, mp 47-49°C.

- 5 Step B: Preparation of 2-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-5-methyl-1H-pyrrol-1-yl]]-1-cyanoethane.


A solution of 1-(4-fluorophenyl)-5-methyl-1,4-hexanedione (36.5 g, 164 mmol), 3-aminopropionitrile  
16 .1/2 fumarate (23.1g, 180.4 mmol), and p-toluenesulfonic acid (0.1 g) in 250 ml of glacial acetic acid was stirred and heated under reflux under nitrogen for six hours. After cooling to room temperature, the mixture was poured into 500 ml of ice-water and the water  
15 suspension which resulted was extracted with two 600-ml portions of ether. The combined ether extract was washed successively with two 200-ml portions of water, three 200-ml portions of sodium bicarbonate, and a 200-ml portion of brine and then dried over anhydrous magnesium  
20 sulfate.

The ether was removed, and the liquid which remained was flash chromatographed on silica gel, eluting with 10:1 (volume/volume) hexane-ethyl acetate to yield 34.8 g of oily 2-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]]-1-cyanoethane which solidified upon  
25 standing.

Recrystallization from isopropyl ether provided analytical material of melting point 78-80°C.

Anal. Calcd. for  $C_{16}H_{17}FN_2$ :

30 C, 74.97%; H, 6.69%; N, 10.93%  
Found: C, 75.18%; H, 6.64%; N, 10.93%.



Step C: Preparation of 3-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]]-1-propanal.

To a stirred solution of 2-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]]-1-cyanoethane (34.8 g, 135.8 mmol) in 300 ml of dichloromethane at ambient temperature under nitrogen was added dropwise over 30 minutes 156.2 ml of a 1.0 M solution of diisobutyl-aluminum ("DiBAL") in dichloromethane. The resulting mixture was stirred for three hours, after which another 20 ml of 1.0 M DiBAL solution was added. The mixture was stirred overnight at room temperature, after which the excess hydrode was destroyed by cautious addition of methanol. When gas evolution had ceased, the solution was carefully poured into 500 ml of vigorously stirred ice-cold 2 M hydrochloric acid.

The emulsion which resulted was extracted with two 500-ml portions of ether and the combined ether extracts were washed successively with 100 ml of water, two 100-ml portions of sodium bicarbonate solution, and 100 ml of brine, and then dried over anhydrous magnesium sulfate. The ether was removed and the residue was flash chromatographed over silica gel, eluting with 10:1 (volume/volume) hexane-ethyl acetate, yielding pure 3-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]]-1-propanal.

Step D: Preparation of methyl 7-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]]-5-hydroxy-3-oxo-heptanoate.

To a stirred suspension of 2.17 g (98.6 mmol) of hexane-washed sodium hydride in 200 ml of anhydrous tetrahydrofuran, cooled to 0°C under nitrogen, was added dropwise over a period of 30 minutes a solution of 8.9 ml (82.4 mmol) of methyl acetoacetate in 150 ml of anhydrous tetrahydrofuran. When gas evolution had ceased, 39.3 ml of a 2.1 M solution of n-butyl lithium in hexane was added dropwise. The resulting solution was stirred for

- 35 -

30 minutes after which a solution of 19.4 g (74.9 mmol) of 3-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]]-1-propanal in 150 ml of anhydrous tetrahydrofuran was added dropwise over a period of 30 minutes. The solution was stirred for an additional hour before quenching the reaction by the addition of 100 ml of saturated aqueous ammonium chloride solution, followed by 100 ml of 2 M hydrochloric acid solution.

The resulting mixture was partitioned between ether (500 ml) and water (100 ml). The water layer was separated and extracted with 300 ml of ether. The ether extracts were combined and washed with 50 ml of brine and then dried over anhydrous magnesium sulfate. The ether was removed and the residue was flash chromatographed on silica gel, eluting with 5:1 (volume/volume) hexane-ethyl acetate to yield 19.9 g (64%) of methyl 7-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]]-5-hydroxy-3-oxo-heptanoate.

20 Step E: Preparation of trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

Thirty ml of air (syringe) were bubbled through a solution of 58 ml of a 1 M solution of tributylborane in tetrahydrofuran containing 19.9 g (53 mmol) of methyl 7-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]]-5-hydroxy-3-oxo-heptanoate under

nitrogen at room temperature. The solution was then stirred for 18 hours at room temperature and then cooled to -78°C. Sodium borohydride (2.27 g, 60 mmol) was then added in one portion. The mixture was  
 5 stirred for 60 minutes at -78°C and for 90 minutes at 0°C. A mixture of 10 ml water and 10 ml of methanol was carefully added (gas evolution). Sixty ml of 3M sodium hydroxide solution and 30 ml of 30% H<sub>2</sub>O<sub>2</sub> solution were simultaneously added  
 10 to the mixture from separation dropping funnels. The vigorously stirred mixture was held at 0°C for 60 minutes and then at room temperature for two hours.

The mixture was then partitioned between 300 ml of water and 300 ml of ether. The ether layer was  
 15 extracted with 50 ml of 10% sodium hydroxide solution and the water layers were combined, acidified with concentrated hydrochloric acid, and extracted with two 500-ml portions of ethyl acetate. The ethyl acetate extracts were combined, washed  
 20 twice with brine, and dried over anhydrous magnesium sulfate. Removal of the ethyl acetate yielded 12.5 g of an oily acid which was dissolved in 500 ml of toluene and heated to azeotropically remove water. After cooling the solution to room temperature  
 25 and removing the toluene, the residue was flash chromatographed on silica gel, eluting with 2:1 hexane-ethyl acetate (volume/volume) to yield 11 g of a colorless solid. Recrystallization from diisopropyl ether yielded 9.5 g (52%) of trans-6-  
 30 [2-{2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl}]tetrahydro-4-hydroxy-2H-pyran-2-one,  
 mp 104-105°C.

Anal. Calcd. for C<sub>20</sub>H<sub>24</sub>FN<sub>3</sub>O<sub>3</sub>:

C, 70.42; H, 7.00; N, 4.06;

35 Found: C, 70.26; H, 7.33; N, 3.99.

## EXAMPLE 2

Preparation of trans-6-[2-[2-(4-fluorophenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one

- 5        The procedure of Example 1 was employed with the substitution of equimolar amounts of 4-fluorobenzaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure of
- 10    Steps B-E were followed to produce trans-6-[2-[2-(4-fluorophenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

Anal. Calcd. for  $C_{18}H_{20}FNO_3$ :

C, 68.12; H, 6.35; N, 4.41;

- 15    Found: C, 68.39; H, 6.18; N, 4.25.

## EXAMPLE 3

Preparation of trans-6-[2-[2-cyclopropyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one

- 20        The procedure of Example 1 was employed with the substitution of equimolar amounts of 4-fluorobenzaldehyde and 1-cyclopropyl-2-propene-1-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter,
- 25    the procedure of Steps B-E were followed to produce trans-6-[2-[2-cyclopropyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

Anal. Calcd. for  $C_{20}H_{22}FNO_3$ :

- 30        C, 69.96; H, 6.46; N, 4.08;

Found: C, 70.02; H, 6.54; N, 4.01.

- 38 -

## EXAMPLE 4

Preparation of trans-6-[2-[2-(1,1-dimethylethyl)-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one

5       The procedure of Example 1 was employed with the substitution of equimolar amounts of 4-fluorobenzaldehyde and t-butyl vinyl ketone for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure  
10 of Steps B-E were followed to produce trans-6-[2-[2-(1,1-dimethylethyl)-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one, mp 177-178°C.

Anal. Calcd. for  $C_{21}H_{26}FNO_3$ :

15       C, 70.17; H, 7.29; N, 3.90;

Found: C, 70.22; H, 7.50; N, 3.80.

## EXAMPLE 5

Preparation of trans-6-[2-(5-phenyl-2-methyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one

20       The procedure of Example 1 was employed with the substitution of equimolar amounts of benzaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure of Steps B-E  
25 were followed to produce trans-6-[2-(5-phenyl-2-methyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one, mp 95-96°C.

Anal. Calcd. for  $C_{19}H_{23}NO_4$ :

C, 69.28; H, 7.04; N, 4.25;

30 Found: C, 68.93; H, 7.00; N, 4.10.

## EXAMPLE 6

Preparation of trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one

- 5        The procedure of Example 1 was employed with the substitution of equimolar amounts of 2-methoxybenzaldehyde and methyl vinyl ketone for the 4-(fluorobenzaldehyde and isopropyl vinyl ketone in Alternate Step A of Example 1. Thereafter, the
- 10      procedure of Steps B-E were followed to produce trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one, mp 112.5-113.5°C.

Anal. Calcd. for  $C_{19}H_{23}NO_4$ :

- 15            C, 69.28; H, 7.04; N, 4.25;  
Found: C, 69.04; H, 7.22; N, 4.17.

## EXAMPLE 7

Preparation of trans-tetrahydro-4-hydroxy-6-[2-[2-(4-methoxyphenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one

- 20        The procedure of Example 1 was employed with the substitution of equimolar amounts of 4-methoxybenzaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in
- 25      Step A of Example 1. Thereafter, the procedure of Steps B-E were followed to produce trans-tetrahydro-4-hydroxy-6-[2-[2-(4-methoxyphenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one, mp 95-95°C.

Anal. Calcd. for  $C_{19}H_{23}NO_4$ :

- 30            C, 69.28; H, 7.04; N, 4.25;  
Found: C, 68.93; H, 7.00; N, 4.10.

## EXAMPLE 8

0179559

Preparation of trans-6-[2-(2-cyclohexyl-5-methyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one

The procedure of Example 1 was employed with the substitution of equimolar amounts of cyclohexanecarboxaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure of Steps B-E were followed to produce trans-6-[2-(2-cyclohexyl-5-methyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one, mp 129-130°C.

Anal. Calcd. for  $C_{18}H_{27}NO_3$ :

C, 70.79; H, 8.91; N, 4.59;

Found: C, 71.11; H, 8.71; N, 4.47.

15

## EXAMPLE 9

Preparation of trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-[3-(trifluoromethyl)phenyl]-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one

The procedure of Example 1 was employed with the substitution of equimolar amounts of 3-(trifluoromethyl)benzaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure of Steps B-E were followed to produce trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-[3-(trifluoromethyl)phenyl]-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one.

Anal. Calcd. for  $C_{19}H_{20}F_3NO_3$ :

C, 62.12; H, 5.49; N, 3.81;

Found: C, 62.22; H, 5.61; N, 3.73.

---

EXAMPLE 10

0179559

Preparation of trans-6-[2-[2-([1,1'-biphenyl]-4-yl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one

5       The procedure of Example 1 was employed with the substitution of equimolar amounts of 4-phenylbenzaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure  
10 of Steps B-E were followed to produce trans-6-[2-[2-([1,1'-biphenyl]-4-yl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one, mp 104-107°C.

Anal. Calcd. for  $C_{24}H_{25}NO_3$ :

15       C, 76.77; H, 6.71; N, 3.73;

Found: C, 76.66; H, 6.66; N, 3.71.

EXAMPLE 11

Preparation of trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(1-naphthalenyl)-1H-pyrrol-1-yl]-2H-pyran-2-one

20       The procedure of Example 1 was employed with the substitution of equimolar amounts of 1-naphthaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in  
25 Step A of Example 1. Thereafter, the procedure of Steps B-E were followed to produce trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(1-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one, mp 137-138°C.

---

0179559

EXAMPLE 12

Preparation of trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(2-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one

5       The procedure of Example 1 was employed with the substitution of equimolar amounts of 2-naphthaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure  
10 of Steps B-E were followed to produce trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(2-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one, mp 45-50°C.  
Anal. Calcd. for C<sub>22</sub>H<sub>23</sub>NO<sub>3</sub>:

C, 75.62; H, 6.63; N, 4.00;

15 Found: C, 75.12; H, 6.88; N, 3.97.

EXAMPLE 13

Preparation of trans-6-[2-(bicyclo[2.2.1]hept-5-en-2-yl-5-methyl-1H-pyrrol-1-yl)ethyl]-tetrahydro-4-hydroxy-2H-pyran-2-one

20       The procedure of Example 1 was employed with the substitution of equimolar amounts of bicyclo[2.2.1]hept-5-ene-2-carboxaldehyde (mixture of diastereomers) and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde  
25 in Step A of Example 1. Thereafter, the procedure of Steps B-E were followed to produce trans-6-[2-(2-bicyclo[2.2.1]hept-5-en-2-yl-5-methyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one as a 1:1 mixture of the endo- and exo-isomers at the  
30 norbornene ring, mp 125-126°C.

Anal. Calcd. for C<sub>19</sub>H<sub>25</sub>NO<sub>3</sub>:

C, 72.35; H, 7.99; N, 4.44;

Found: C, 72.11; H, 8.02; N, 4.32.

0179559

EXAMPLE 14

Preparation of trans-6-[2-[2-(diphenylmethyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one

- 5        The procedure of Example 1 was employed with the substitution of equimolar amounts of diphenylacetaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure
- 10 of Steps B-E were followed to produce trans-6-[2-[2-diphenylmethyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one, mp 129-132°C.
- Anal. Calcd. for  $C_{25}H_{27}NO_3$ :  
C, 77.07; H, 6.99; N, 3.60;
- 15 Found: C, 76.85; H, 7.14; N, 3.45.

EXAMPLE 15

Preparation of trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]propyl]tetrahydro-4-hydroxy-2H-pyran-2-one

- 20        The procedure of Example 1 was employed with the substitution in Step B of 2-aminopropanol in place of the ethanolamine. Thereafter, the procedure of Steps C-E were followed to produce trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-
- 25 yl]propyl]tetrahydro-4-hydroxy-2H-pyran-2-one, mp 167-169°C.
- Anal. Calcd. for  $C_{21}H_{26}FNO_3$ :  
C, 70.17; H, 7.29; N, 3.90;
- Found: C, 70.06; H, 7.36; N, 3.82.
-

EXAMPLE 16

Preparation of trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxyphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl-ethyl]-2H-pyran-2-one

5        The procedure of Example 1 was employed with the substitution of equimolar amounts of 2-methoxybenzaldehyde and 3-butene-2-one for the 1-(4-fluorophenyl)-2-propene-1-one and isobutyraldehyde in Step A of Example 1. Thereafter, the procedure of Steps B-E  
10        were followed to produce trans-tetrahydro-4-hydroxy-6-2-[2-(2-methoxyphenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl-ethyl]-2H-pyran-2-one.

Anal. Calcd. for  $C_{21}H_{27}NO_3$ :

C, 70.56; H, 7.61; N, 3.92;

15        Found: C, 70.43; H, 7.66; N, 3.73.

---

## EXAMPLE 17

## Method 1

Step A: Preparation of 6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-tert-butyl dimethylsilyloxy-, trans-2H-pyran-2-one.

To a solution of 6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]-tetrahydro-4-hydroxy-trans-2H-pyran-2-one (0.52 g, 1.5  $\mu$ moles) and tert-butyldimethylchloro silane (0.27 g, 1.8  $\mu$ moles) in 5 ml of dry DMF was added imidazole (0.31 g, 4.5  $\mu$ moles) in one portion. The solution was stirred overnight at room temperature before partitioning between hexane (100 ml) and water (50 ml). The aqueous layer was extracted with two 50 ml portions of hexane. The combined hexane extracts were washed with H<sub>2</sub>O (2 x 25 ml), brine (25 ml), and dried (MgSO<sub>4</sub>). Filtration through silica gel and concentration provided 0.7 g (100%) of the title compound. 90 MHz NMR (CDCl<sub>3</sub>)  $\delta$  0.10 (s, 6H), 0.90 (s, 9H), 1.30 (d, J=Hz 6H), 1.4-1.8 (m, 4H), 2.48 (m, 2H), 2.95 (m, 1H), 3.9-4.3 (m, 3H), 5.85 (d, J=2Hz 1H), 6.02 (d, J=2Hz, 1H), 6.8-7.3 (m, 4H).

Step B: Preparation of 6-[2-[2-(4-fluorophenyl)-3,4-dichloro-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-trans-2H-pyran-2-one.

N-Chlorosuccinimide (6.48  $\mu$ moles, 0.87 g) was added in one portion to a stirred solution of 6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-tert-butyldimethylsilyloxy-trans-2H-pyran-2-one (1.49 g, 3.24  $\mu$ moles) in dry DMF (10 ml) cooled to 0°C under dry nitrogen. The solution was stirred for one hour at 0°C then warmed to room temperature for three hours. It was then diluted with water (50 ml) and extracted with ether (2 x 1000 ml). The ether extracts were diluted with 100 ml of hexane and washed with water

- (50 ml), 10% aq.  $\text{NaHCO}_3$  (50 ml), 10% aq.  $\text{NaHSO}_3$  (50 ml), brine (50 ml), and dried ( $\text{MgSO}_4$ ). The crude product which remained after filtration and concentration was dissolved in tetrahydrofuran (15 ml) and treated with
- 5 glacial acetic acid (0.75 ml, 13 mmol) and tetrabutyl ammonium fluoride (9.72 ml of 1 M THF solution). The solution was stirred for five hours, diluted with ethyl acetate (100 ml) and washed with saturated aq.  $\text{NaHCO}_3$  (2 x 50 ml), brine (25 ml), and dried ( $\text{MgSO}_4$ ).
- 10 The residue which remained after filtration and concentration was flash chromatographed on silica gel eluting with 2:1 hexane-ethyl acetate. This provided 0.50 g (35%) of pure lactone. Recrystallization from ether-hexane provided colorless crystals mp 129-131°C.
- 15 Anal. Calcd. for  $\text{C}_{20}\text{H}_{22}\text{FCl}_2\text{NO}_3$ :  
C, 57.98; H, 5.35; N, 3.38;  
Found: C, 58.24; H, 5.24; N, 3.39.  
IR (KBr)  $\nu$  3550, 2990, 1711, 1518, 12225, 1160, 1055, 851, 816  $\text{cm}^{-1}$  200 MHz NMR ( $\text{CDCl}_3$ )  $\delta$  1.44 (d, J=7Hz, 6H),  
20 1.8 (m, 4H), 2.12 (d, J=3Hz, 1H, -OH), 2.55 (m, 2H), 3.10 (M, 1H), 4.0 (M, 2H), 4.30 (M, 1H), 4.45 (M, 1H), 7.0-7.4 (M, 4H).

## Method 2

- Step A: Preparation of 2-(4-fluorophenyl)-5-(1-methylethyl)-3,4-dichloro-1H-pyrrole-1-propanenitrile.
- 25 N-Chlorosuccinimide (practical, 105 g, 786.5 mmol) was added in one portion to a stirred solution of 2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrole-1-propanenitrile
- 30 (84 g, 327.7 mmol) in 500 ml of dry dimethylformamide cooled to 0°C under nitrogen. After stirring for 60 minutes at 0°C and 90 minutes at 25°C, a further 8 g (60 mmol) of N-chlorosuccinimide were added. The solution was stirred a further 60 minutes before pouring into
- 35 ether (3 liters) and washing with  $\text{H}_2\text{O}$  (3 x 500 ml), 10% aq.  $\text{NaHSO}_3$  (300 ml),  $\text{H}_2\text{O}$  (300 ml), brine, and dried ( $\text{MgSO}_4$ ). Flash chromatography on silica gel eluting with

10:1 hexane-ethyl acetate provided an oil which solidified on standing. Recrystallization from isopropyl ether-hexane provided 96 g of colorless crystals mp 80-82°C.

5 Anal. Calcd. for  $C_{16}H_{15}Cl_2FN_2$ :

C, 59.09; H, 4.65; N, 8.61;

Found: C, 59.01; H, 4.56; N, 8.59.

IR (KBr) 2933, 2249, 1520, 1490; 1344, 1315, 1218, 848, 524  $cm^{-1}$ . 100 MHz NMR ( $CDCl_3$ )  $\delta$  1.42 (d, J=7Hz,

10 6H), 2.33 (t, J=7Hz, 2H), 3.0 (sptet, J=7Hz, 1H), 4.05 (t, J=7Hz, 2H), 70-74 (m, 4H).

Employing the product of this step in the process described above in Step C of Example 1, provided 6-[2-[2-(4-fluorophenyl)-3,4-dichloro-5-(1-methylethyl)-1H-pyrrol-1-yl]-ethyl]tetrahydro-4-  
15 hydroxy-trans-2H-pyran-2-one.

#### EXAMPLE 18

20 Preparation of 6-[2-[2-(4-fluorophenyl)-3,4 dibromo-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-trans-2H-pyran-2-one.

Substitution of N-Bromosuccinimide for N-Chlorosuccinimide in Step B of Method 1, Example 17 provided  
25 a corresponding amount of the title compound mp 143°C.

Anal. Calcd. for  $C_{20}H_{22}FBr_2NO_3$ :

C, 47.74; H, 4.41; N, 2.78; Br, 31.76; F, 3.77.

Found: C, 47.52; H, 4.34; N, 2.84. Br, 31.75; F, 3.72.

30 IR (KBr) 3350, 2966, 1711, 1510, 1484, 1225, 1072, 847, 820  $cm^{-1}$ . 200 MHz NMR ( $CDCl_3$ )  $\delta$  1.40 (d, J=7Hz, 6H), 1.5-1.8 (m, 4H), 1.94 (brs, 1H, -OH), 2.58 (m, 2H), 3.13 (m, 1H), 4.0 (m, 2H), 4.31 (m, 1H), 4.47 (m, 1H), 7.0-7.3 (m, 4H).

## EXAMPLE 19

Step A: Preparation of ethyl-2-(1-(1-oxo-2,2,2-trifluoroethyl))-4-oxo-4-(4-fluorophenyl)-butyrate

A solution of ethyl 1,1,1,-trifluoroacetoacetate (14.6 ml, 0.1 mole) in dry DMF (100 ml) was added dropwise to a 0°C suspension of hexane washed sodium hydride (0.106 mole) in 50 ml of dry DMF under nitrogen. When gas evolution was complete, a solution of  $\alpha$ -bromo-4'-fluoroacetophenone (0.1 mole,) prepared as in J. Org. Chem. 29, 3459 (1964)) in 100 ml of dry DMF was added dropwise over 30 minutes. The mixture was allowed to warm slowly to 25°C overnight. It was then quenched by addition of 6 N HCl, poured into H<sub>2</sub>O (1 liter) and extracted with ether (2 x 500 ml). The combined ether extracts were washed with H<sub>2</sub>O (2 x 100 ml), brine (100 ml), and dried (MgSO<sub>4</sub>). Flash chromatography on silica gel eluting with 5:1 hexane-ethylacetate provided 7 g of the title compound. IR (film) 3380, 1768, 1744, 1688, 1601, 1511, 1413, 1293, 1263, 1238, 1212, 1160, 1100, 1004, 841 cm<sup>-1</sup>. 200 MHz NMR (CDCl<sub>3</sub>)  $\delta$  1.29 (t, J=7Hz, 3H), 3.75 (m, 2H), 4.26 (q, J=7Hz, 2H), 4.55 (dd, J=4.7, 9.6Hz, 1H), 7.21 (m, 2H), 8.02 (m, 2H)

Step B: Preparation of 2-(4-fluorophenyl)-5-trifluoromethyl-1H-pyrrole-1-propanenitrile.

A solution of ethyl-2-(1-(1-oxo-2,2,2-trifluoroethyl))-4-oxo-4-(4-fluorophenyl)-butyrate (5 g, 15.6  $\mu$ moles) in 110 ml of 5:5:1 acetic acid-water-conc. sulfuric acid was stirred and heated at reflux for four hours. The cooled solution was carefully poured into 400 ml of saturated aq. bicarbonate which was then extracted with ether (2 x 300 ml). The combined ether extracts were washed with saturated aq. bicarbonate (2 x 50 ml), brine (50 ml), and dried (MgSO<sub>4</sub>). The crude

- diketone which remained after filtration and concentration (3 g) was dissolved in 20 ml of glacial acetic acid and 2 g (18 mmoles) of 3-aminopropanenitrile-1/2-fumarate were added. The solution was stirred and
- 5 heated at reflux for five hours. The cooled solution was poured into 200 ml of saturated aq. bicarbonate and extracted with ether (2 x 200 ml). The combined ether extracts were washed with H<sub>2</sub>O (2 x 50 ml), brine (50 ml), and dried (MgSO<sub>4</sub>). Flash chromatography of the
- 10 residue which remained after filtration and concentration provided 1.2 g (27%) of the title compound.
- IR (CDCl<sub>3</sub>) 2258, 1611, 1570, 1478, 1337, 1172, 1106, 1064, 844 cm<sup>-1</sup>. 200 MHz NMR (CDCl<sub>3</sub>)  $\delta$  2.51 (t, J=7.3Hz, 2H), 4.30 (t, J=7Hz, 2H), 6.16 (d, J=3.8Hz,
- 15 1H), 6.67 (d, J=3.8Hz, 1H), 7.1-7.5 (m, 4H). Mass spectrum M/e 282, 263, 242, 229, 173.
- Preparation of 6-[2-[2-trifluoromethyl]-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-trans-2H-pyran-2-one.
- 20 Substitution of 2-(4-fluorophenyl)-5-trifluoromethyl-1H-pyrrole-1-propanenitrile for 2-(4-fluorophenyl)-3,4-dichloro-5-(1-methylethyl)-1H-pyrrole-1-propanenitrile in Step C of Example 1 and following the procedures of Step C, D, and E resulted in a corresponding amount of
- 25 the title compound as an oil.
- Anal. Calcd. for C<sub>18</sub>H<sub>17</sub>F<sub>4</sub>NO<sub>3</sub>:
- C, 58.22; H, 4.61; N, 3.77.
- Found: C, 58.88; H, 5.07; N, 4.03.
- 30 IR (film) 3440, 2927, 1728, 156, 1477, 1342, 1266, 1230, 1160, 1101, 1060, 843, 782 cm<sup>-1</sup>. 200 MHz NMR (CDCl<sub>3</sub>)  $\delta$  1.3-2.1 (m, 4H), 2.34 (brs, 1H, -OH), 2.55 (m, 2H), 3.9-4.3 (m, 3H), 4.52 (m, 1H), 6.11 (d, J=3.8Hz, 1H), 6.61 (dd, J=0.8, 3.8Hz, 1H), 7.0-7.4 (m, 4H).

## EXAMPLE 20

Preparation of (±)-N-(4-fluorobenzoyl)-N-[2-(2-ethyl)-1,3-dioxolanyl]valine.

5        A solution of the methyl-2-bromo-3-methyl butyrate (4.6 g, 23.6 mmol), 2-(1-(2-aminoethyl))-1,3-dioxolane (2.93 g, 25 mmol) and triethylamine (3.5 ml, 25 mmol) was stirred and heated in 25 ml of refluxing acetonitrile for 20 hours. The cooled solution was poured into ether  
10    (500 ml) and extracted 2M HCl (2 x 50 ml). The aqueous layer was made alkaline with 25% aq. NaOH and extracted with ethyl acetate (2 x 100 ml). The combined ethyl acetate extracts were washed with brine and dried (MgSO<sub>4</sub>). Filtration and concentration provided 3 g of the title  
15    compound as liquid. 90 MHz NMR (CDCl<sub>3</sub>) δ 0.9.3 (d, J=7Hz, 6H), 1.70 (brs, 1H, -NH), 1.86 (m, 2H), 2.60 (m, 3H) 2.94 (d, J=6Hz, 1H), 3.68 (s, 3H), 3.85 (m, 4H), 4.89 (t, J=4Hz, 1H).

20    Preparation of ±-Methyl-N-(4-fluorobenzoyl)-N-[2-(2-ethyl)-1,3 dioxolanyl]valine.

      To a stirred solution of Methyl-N-[2-(2-ethyl)-1,3-dioxolanyl]valine (3 g, 13 mmol) and triethylamine (3.6 ml, 26 mmol) in 20 ml of dichloromethane (CH<sub>2</sub>CL<sub>2</sub>) cooled to 0°C was added a solution of 4-fluorobenzoyl  
25    chloride (1.65 ml, 14 mmol) in 10 ml of (CH<sub>2</sub>CL<sub>2</sub>). The solution was stirred 60 minutes at 0°C and 60 minutes at room temperature. It was then poured into ether and washed with water, saturated aq. bicarbonate, brine, and  
30    dried (MgSO<sub>4</sub>). Flash chromatography on silica gel eluting with 1:1 hexane-ethyl acetate provided 3 g of the title compound. 90 MHz NMR (CDCl<sub>3</sub>) δ 0.90, (brd, J=7Hz, 6H), 1.8-2.5 (m, 3H), 3.45 (br, dd, J=6, 8Hz, 1H), 3.72 (s, 3H), 3.80 (m, 6H), 4.80 (m, 1H),  
35    6.9-7.5 (m, 4H).

Preparation of  $\alpha$ -N-(4-fluorobenzoyl)-N-[2-(2-ethyl)-1,3-dioxolanyl]valine.

A solution of the methyl ester prepared above (1.9 g, 2.83 mmol) and NaOH (0.4 g, 10 mmol) in 10 ml of 4:1 CH<sub>3</sub>OH-H<sub>2</sub>O was stirred and heated at reflux for three hours. The cooled solution was diluted with water and extracted with ether. The aqueous layer was acidified with 6M HCl and extracted with ethyl acetate. (2x). The combined ethyl acetate extracts were washed with brine and dried (MgSO<sub>4</sub>). Filtration and concentration provided 0.96 g (2.8 mmol) of acid. 90MHz NMR (CDCl<sub>3</sub>)  $\delta$  0.85 (m, 6H), 1.8 (m, 2H), 2.5 (m, 1H) 3.3-3.9 (m, 7H), 4.6 (m, 1H) 6.8-7.4 (m, 4H).

Preparation of dimethyl-1-[2-(2-ethyl)-1,3-dioxolanyl]dioxolanyl]-2-(4-fluorophenyl)-5-(1-methyl-ethyl)-1H-pyrrole-3,4-dicarboxylate

Dimethyl acetylene dicarboxylate (1.3 ml, 10.6 mmol) was added to a 25°C solution of (+)-N-(4-fluorobenzoyl)-N-[2-(2-ethyl)-1,3-dioxolanyl]valine (1.8 g, 5.28 mmol) dissolved in 10 ml of acetic anhydride. The evolution of carbon dioxide began immediately. The solution was stirred a further two hours, concentrated to remove excess acetylene and solvent, then filtered through silica gel. This provided 2 g of pyrrole as a solid which was recrystallized from isopropyl ether-hexane mp 143-146°C.

Anal. Calcd. for C<sub>22</sub>H<sub>26</sub>FNO<sub>6</sub>

C, 62.55; H, 6.20; N, 3.31.

Found: C, 62.84; H, 6.23; N, 3.30.

IR (KBr) 1719, 1449, 1241, 1209, 1178, 945 cm<sup>-1</sup>.

200 MHz NMR (CDCl<sub>3</sub>)  $\delta$  1.35 (d, J=7Hz, 6H), 1.80 (m, 2H), 3.18 (Septet, J=7Hz, 1H), 3.56 (s, 3H), 1H), 3.7-4.0 (m, 6H), 3.83 (s, 3H), 4.64 (t, J=4Hz, 1H), 7-7.3 (m, 4H).

Preparation of Dimethyl-1-(1-(3-oxopropyl))-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrole-3,4-dicarboxylate

A solution of dimethyl-1-[2-(2-ethyl)-1,3-dioxolanyl]-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrole-3,4-dicarboxylate (0.5 g, 1.18 mmol) and p-toluenesulfonic acid (0.23 g, 1.2 mmol) in 12 ml of 5:1 acetone-water was stirred and heated at reflux for 48 hours. The cooled solution was concentrated, diluted with ether (200 ml), washed with saturated aq. bicarbonate (2 x 50 ml), brine (50 ml), and dried (MgSO<sub>4</sub>). Flash chromatography on silica gel eluting with 4:1 hexane-ethyl acetate provided 0.4 g of pure aldehyde. 90 MHz NMR (CDCl<sub>3</sub>)  $\delta$  1.35 (d, J=7Hz, 6H), 2.61 (t, J=7Hz, 2H), 3.18 (septet, J=7Hz, 1H), 3.53 (s, 3H), 3.81 (s, 3H), 4.03 (t, J=7Hz, 2H), 6.9-7.3 (m, 4H), 9.45 (s, 1H).

Preparation of Dimethyl-2-(4-Fluorophenyl)-5-(1-methylethyl)-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3,4-dicarboxylate.

Substitution of dimethyl-1-(1-(3-oxopropyl))-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrole-3,4-dicarboxylate for 2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrole-1-propanal in Step C of Example 1 and following the procedures of Steps C, D, and E provided a corresponding amount of the title compound mp 167-170°C. Anal. Calcd. for C<sub>24</sub>H<sub>28</sub>FNO<sub>7</sub>


C, 62.47; H, 6.12; N, 3.04.

Found: C, 62.32; H, 5.87; N, 2.99.

IR (KBr) 2450, 2980, 1719, 1499, 1225, 1174, 1074, 811 cm<sup>-1</sup>. 200 MHz NMR (CDCl<sub>3</sub>)  $\delta$  1.34 (d, J=7Hz, 6H), 1.57 (m, 4H), 2.40 (d, J=3Hz, 1H), 2.56 (m, 2H), 3.16 (septet, J=7Hz, 1H), 3.55 (s, 3H), 3.83 (s, 3H), 4.0 (m, 2H), 4.26 (m, 1H), 4.44 (m, 1H), 4.44 (m, 1H), 7.1-7.3 (m, 4H).

CLAIMS: (for BE, CH, DE, FR, GB, IT, LI, LU, NL, ~~DE~~ 79559)

1. A compound having the structural formula I:



wherein X is

-CH<sub>2</sub>-,  
 -CH<sub>2</sub>CH<sub>2</sub>-, or  
 -CH(CH<sub>3</sub>)CH<sub>2</sub>-;

R<sub>1</sub> is

1-naphthyl,  
 2-naphthyl,  
 cyclohexyl,  
 norbornenyl,  
 phenyl,  
 phenyl substituted by  
 fluorine,  
 chlorine,  
 hydroxy,  
 trifluoromethyl,  
 alkyl of from one to four carbon atoms,  
 alkoxy of from one to four carbon atoms, or  
 alkanoyloxy of from two to eight carbon  
 atoms,  
 2-, 3-, or 4-pyridinyl,  
 2-, 3-, or 4-pyridinyl-N-oxide, or




-54-

where  $R_5$  is alkyl of from one to four carbon atoms and  $\text{hal}^-$  is chloride, bromide, or iodide;


$R_2$  and  $R_3$  are independently

- 5 hydrogen,  
chlorine,  
bromine,  
cyano,  
trifluoromethyl,  
10 phenyl,  
alkyl of from one to four carbon atoms,  
carboalkoxy of from two to eight carbon atoms,  
 $-\text{CH}_2\text{OR}_6$  where  $R_6$  is  
hydrogen,  
15 alkanoyl of from one to six carbon atoms,  
 $-\text{CH}_2\text{OCONHR}_7$  where  $R_7$  is  
alkyl of from one to six carbon atoms,  
phenyl,  
phenyl substituted with  
20 chlorine,  
bromine, or  
alkyl of from one to four carbon atoms;  
or when taken together with the carbon atoms to  
25 which they are attached,  $R_2$  and  $R_3$  form a  
ring denoted by




where  $n$  is three or four,

a ring denoted by



a ring denoted by



5

where  $R_8$  is

hydrogen,


alkyl of from one to six carbon  
atoms,

10

phenyl, or

benzyl;

or a ring denoted by



15

where  $R_9$  and  $R_{10}$  are

20

hydrogen,

alkyl of from one to four carbon  
atoms, or

benzyl;

25

$R_4$  is

alkyl of from one to four carbon atoms,

cyclopropyl,

cyclobutyl, or

trifluoromethyl;


30

or a corresponding lactone ring-opened dihydroxy  
acid derived therefrom, or a pharmaceutically  
acceptable salt thereof.

35

2. A compound in accordance with Claim 1, wherein

X is



5

$R_1$  is as defined in Claim 1;

$R_2$  and  $R_3$  are independently

hydrogen,

10


chlorine, or

bromine; and

$R_4$  is as defined in Claim 1.

15 3. A compound in accordance with Claim 1, wherein

X is



20

$R_1$  is

phenyl,

phenyl substituted by

fluorine,

chlorine,

25

hydroxy,

trifluoromethyl,

alkyl of from one to four carbon atoms,

alkoxy of from one to four carbon atoms,

alkanoyloxy of from two to eight carbon


30

atoms,

2-, 3-, or 4-pyridinyl,

2-, 3-, or 4-pyridinyl-N-oxide, or

35



where  $R_5$  is alkyl of from one to four

- 57 -

carbon atoms and  $\text{hal}^-$  is chloride,  
bromide, or iodide;

5  $R_2$  and  $R_3$  are independently  
hydrogen,  
chlorine,  
bromine; and

10  $R_4$  is  
alkyl of from one to four carbon atoms, or  
trifluoromethyl.

4. A compound in accordance with Claim 1, wherein

15 X is  $-\text{CH}_2\text{CH}_2-$ ;

$R_1$  is  
phenyl, or  
phenyl substituted by  
20 fluorine,  
chlorine,  
hydroxy,  
trifluoromethyl,  
alkoxy of from one to four carbon atoms,  
25 alkanoyloxy of from two to eight carbon  
atoms;

$R_2$  and  $R_3$  are independently  
hydrogen,  
30 chlorine, or  
bromine; and

$R_4$  is isopropyl or trifluoromethyl.

35

5. A compound in accordance with Claim 1, wherein

X is  $-\text{CH}_2\text{CH}_2-$ ;

5  $\text{R}_1$  is

phenyl,

phenyl substituted by

fluorine,

chlorine,

10

trifluoromethyl,

alkyl of from one to four carbon atoms,

alkoxy of from one to four carbon atoms,

alkanoyloxy of from two to eight carbon

atoms,

15

1-naphthyl,

2-naphthyl;

$\text{R}_2$  and  $\text{R}_3$  are independently

hydrogen,

20

chlorine,

bromine,

cyano,

trifluoromethyl,

phenyl,

25

alkyl of from one to four carbon atoms,

carboalkoxy of from two to eight carbon atoms,

$-\text{CH}_2\text{OR}_6$  where  $\text{R}_6$  is

hydrogen or alkanoyl of from one to six

carbon atoms,

30

$-\text{CH}_2\text{CONHR}_7$  where  $\text{R}_7$  is

alkyl of from one to six carbon atoms,

phenyl, or

phenyl substituted with

chlorine,

35


bromine, or

alkyl of from one to four carbon

atoms;


or, when taken together with the carbon atoms to which they are attached,  $R_2$  and  $R_3$  form a ring denoted by

5




where  $n$  is three or four;  
a ring denoted by

10



a ring denoted by

15



where  $R_8$  is

20

hydrogen,


alkyl of from one to four carbon  
atoms,

phenyl, or

benzyl, or

25

a ring denoted by



30

where  $R_9$  and  $R_{10}$  are  
hydrogen,

alkyl of from one to four carbon  
atoms, or

benzyl; and

35

$R_4$  is

alkyl of from one to four carbon atoms,  
cyclopropyl,

cyclobutyl, or  
trifluoromethyl.

6. A compound in accordance with Claim 1, wherein

5 X is  $-\text{CH}_2\text{CH}_2-$ ;


R<sub>1</sub> is

phenyl,  
phenyl substituted by  
10 fluorine,  
chlorine,  
trifluoromethyl,  
alkyl of from one to four carbon atoms,  
alkoxy of from one to four carbon atoms,  
15 or alkanoyloxy of from two to eight carbon  
atoms;

R<sub>2</sub> and R<sub>3</sub> are independently


hydrogen,  
20 chlorine,  
bromine,  
phenyl,  
carboalkoxy of from two to eight carbon atoms,  
or, when taken together with the carbon  
25 atoms to which they are attached, R<sub>2</sub> and R<sub>3</sub>  
form a ring  
denoted by

30



where n is three or four;  
a ring denoted by


35



where  $R_8$  is  
hydrogen, or  
alkyl of from one to four carbon  
atoms; or

0179559

5 a ring denoted by



10 where  $R_9$  and  $R_{10}$  are  
hydrogen or alkyl of from one to  
four carbon atoms; and

$R_4$  is  
15 alkyl of from one to four carbon atoms, or  
trifluoromethyl.

7. A compound in accordance with Claim 1, wherein


20 X is  $-\text{CH}_2\text{CH}_2-$ ,

$R_1$  is  
phenyl, or  
phenyl substituted by  
25 fluorine,  
chlorine,  
trifluoromethyl,  
alkyl of from one to four carbon  
atoms,  
30 alkoxy of from one to four carbon  
atoms, or  
alkanoyloxy of from two to eight carbon  
atoms;

35  $R_2$  and  $R_3$  are independently  
carboalkoxy of from two to eight carbon  
atoms or,

0179559

when taken together with the carbon atoms to which they are attached form a ring denoted by




wherein R<sub>8</sub> is hydrogen or alkyl of from one to four carbon atoms; and R<sub>4</sub> is isopropyl or trifluoromethyl.

8. A compound in accordance with Claim 1, selected from the group consisting of trans-6-[2-[3,4-dichloro-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- trans-6-2-[3,4-dibromo-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- trans-6-[2-[2-(4-fluorophenyl)-5-(trifluoromethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- trans-dimethyl 2-(4-fluorophenyl)-5-(1-methylethyl)-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3,4-dicarboxylate;
- trans-6-[2-[2-(4-fluorophenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- trans-6-[2-[2-cyclopropyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- trans-6-[2-[2-(1,1-dimethylethyl)-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;

trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxy-phenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-2-one;  
trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxy-phenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;  
trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(1-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;  
trans-6-[2-(2-bicyclo[2.2.1]hept-5-en-2-yl-5-methyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one; and  
trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylphenyl)-1H-pyrrol-1-yl]propyl]tetrahydro-4-hydroxy-2H-pyran-2-one.


9. A method of preparing a compound having the structural formula




wherein X, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> are as defined in Claim 1, said method comprising the steps of:

-64-

(a) first reacting a substituted [(pyrrol-1-yl)-alkyl]aldehyde compound of Formula III




where X, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are as defined above, with the alkali metal salt of the dianion of methyl acetoacetate to form a compound of structural Formula IV



where X, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are as defined above, then successively

(b) reducing Compound IV with a trialkylborane and sodium borohydride, and

(c) oxidizing with alkaline hydrogen peroxide to produce an acid compound of Formula V,



35

(d) cyclizing, if desired, the acid compound of Formula V to a lactone compound of Formula I by heating in an inert solvent or, alternatively converting, if desired, the acid compound of Formula V to a pharmaceutically acceptable salt.

5  
10 A pharmaceutical composition, useful as a hypocholesterolemic agent, comprising a hypocholesterolemic effective amount of a compound in accordance with any one of Claims 1 to 8 in combination with a pharmaceutically acceptable carrier or diluent.

11. For use in a method of treatment in which cholesterol biosynthesis in a patient is inhibited, a compound in accordance with any one of Claims 1 to 8 or a pharmaceutical composition in accordance with Claim 10.

20


25

30

35

**CLAIMS:** (for AT):

1. A process for preparing a compound having the structural formula 1:



wherein X is

- CH<sub>2</sub>-,
- CH<sub>2</sub>CH<sub>2</sub>-, or
- CH(CH<sub>3</sub>)CH<sub>2</sub>-;

R<sub>1</sub> is

- 1-naphthyl,
- 2-naphthyl,
- cyclohexyl,
- norbornenyl,
- phenyl,
- phenyl substituted by
- fluorine,
- chlorine,
- hydroxy,
- trifluoromethyl,
- alkyl of from one to four carbon atoms,
- alkoxy of from one to four carbon atoms, or
- alkanoyloxy of from two to eight carbon
- atoms,
- 2-, 3-, or 4-pyridinyl,
- 2-, 3-, or 4-pyridinyl-N-oxide, or




-67-

where  $R_5$  is alkyl of from one to four carbon atoms and  $\text{hal}^-$  is chloride, bromide, or iodide;

$R_2$  and  $R_3$  are independently


- 5 hydrogen,  
chlorine,  
bromine,  
cyano,  
trifluoromethyl,  
10 phenyl,  
alkyl of from one to four carbon atoms,  
carboalkoxy of from two to eight carbon atoms,  
 $-\text{CH}_2\text{OR}_6$  where  $R_6$  is  
hydrogen,  
15 alkanoyl of from one to six carbon atoms,  
 $-\text{CH}_2\text{OCONHR}_7$  where  $R_7$  is  
alkyl of from one to six carbon atoms,  
phenyl,  
phenyl substituted with  
20 chlorine,  
bromine, or  
alkyl of from one to four carbon atoms;  
or when taken together with the carbon atoms to  
25 which they are attached,  $R_2$  and  $R_3$  form a  
ring denoted by



30

where  $n$  is three or four,


a ring denoted by



35

a ring denoted by

0179559




5

where  $R_8$  is  
hydrogen,  
alkyl of from one to six carbon  
atoms,  
phenyl, or  
benzyl;

10

or a ring denoted by



15

where  $R_9$  and  $R_{10}$  are  
hydrogen,  
alkyl of from one to four carbon  
atoms, or  
benzyl;

20

25


$R_4$  is  
alkyl of from one to four carbon atoms,  
cyclopropyl,  
cyclobutyl, or  
trifluoromethyl;

30


or a corresponding lactone ring-opened dihydroxy  
acid derived therefrom, or a pharmaceutically  
acceptable salt thereof;  
which process comprises:

35

(a) first reacting a substituted [(pyrrol-1-yl)-alkyl]aldehyde compound of Formula III




10 where X, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are as defined above, with the alkali metal salt of the dianion of methyl acetoacetate to form a compound of structural Formula IV



where X, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are as defined above, then successively

20 (b) reducing Compound IV with a trialkylborane and sodium borohydride, and

(c) oxidizing with alkaline hydrogen peroxide to produce an acid compound of Formula V,



and finally

35

(d) cyclizing, if desired, the acid compound of  
Formula V to a lactone compound of Formula I by  
heating in an inert solvent or, alternatively  
converting, if desired, the acid compound of  
Formula V to a pharmaceutically acceptable salt.

5

10


15

20

25


30

35



2. A process in accordance with Claim 1, wherein

X is



5

$R_1$  is as defined in Claim 1;

$R_2$  and  $R_3$  are independently  
hydrogen,


10

chlorine, or  
bromine; and

$R_4$  is as defined in Claim 1.

15 3. A process in accordance with Claim 1, wherein

X is



20

$R_1$  is

phenyl,

phenyl substituted by

fluorine,

chlorine,

25

hydroxy,

trifluoromethyl,

alkyl of from one to four carbon atoms,

alkoxy of from one to four carbon atoms,


alkanoyloxy of from two to eight carbon  
atoms,

30

2-, 3-, or 4-pyridinyl,

2-, 3-, or 4-pyridinyl-N-oxide, or

35



where  $R_5$  is alkyl of from one to four

carbon atoms and  $\text{hal}^-$  is chloride,  
bromide, or iodide;

5  $R_2$  and  $R_3$  are independently  
hydrogen,  
chlorine,  
bromine; and

10  $R_4$  is  
alkyl of from one to four carbon atoms, or  
trifluoromethyl.

4. A process in accordance with Claim 1, wherein

15 X is  $-\text{CH}_2\text{CH}_2-$ ;

$R_1$  is  
phenyl, or  
phenyl substituted by  
20 fluorine,  
chlorine,  
hydroxy,  
trifluoromethyl,  
alkoxy of from one to four carbon atoms,  
25 alkanoyloxy of from two to eight carbon  
atoms;

$R_2$  and  $R_3$  are independently  
hydrogen,  
30 chlorine, or  
bromine; and

$R_4$  is isopropyl or trifluoromethyl.

35

5. A process in accordance with Claim 1, wherein

X is  $-\text{CH}_2\text{CH}_2-$ ;

5  $R_1$  is

phenyl,

phenyl substituted by

fluorine,

chlorine,

10 trifluoromethyl,

alkyl of from one to four carbon atoms,

alkoxy of from one to four carbon atoms,

alkanoyloxy of from two to eight carbon  
atoms,

15 1-naphthyl,

2-naphthyl;

$R_2$  and  $R_3$  are independently

hydrogen,

20 chlorine,

bromine,

cyano,

trifluoromethyl,

phenyl,

25 alkyl of from one to four carbon atoms,

carboalkoxy of from two to eight carbon atoms,

$-\text{CH}_2\text{OR}_6$  where  $R_6$  is

hydrogen or alkanoyl of from one to six  
carbon atoms,

30  $-\text{CH}_2\text{OCONHR}_7$  where  $R_7$  is

alkyl of from one to six carbon atoms,

phenyl, or

phenyl substituted with

chlorine,

35 bromine, or

alkyl of from one to four carbon  
atoms;


or, when taken together with the carbon atoms to which they are attached,  $R_2$  and  $R_3$  form a ring denoted by



where  $n$  is three or four;  
a ring denoted by




a ring denoted by



where  $R_8$  is

20 hydrogen,  
alkyl of from one to four carbon  
atoms,  
phenyl, or  
benzyl, or

a ring denoted by



where  $R_9$  and  $R_{10}$  are  
hydrogen,

alkyl of from one to four carbon 78  
atoms, or  
benzyl; and

35  $R_4$  is

alkyl of from one to four carbon atoms,  
cyclopropyl,

cyclobutyl, or  
trifluoromethyl.

0179559

6. A process in accordance with Claim 1, wherein

5 X is  $-\text{CH}_2\text{CH}_2-$ ;

R<sub>1</sub> is

phenyl,  
phenyl substituted by  
10 fluorine,  
chlorine,  
trifluoromethyl,  
alkyl of from one to four carbon atoms,  
alkoxy of from one to four carbon atoms,  
15 or alkanoyloxy of from two to eight carbon  
atoms;

R<sub>2</sub> and R<sub>3</sub> are independently

hydrogen,  
20 chlorine,  
bromine,  
phenyl,  
carboalkoxy of from two to eight carbon atoms,  
or, when taken together with the carbon  
25 atoms to which they are attached, R<sub>2</sub> and R<sub>3</sub>  
form a ring  
denoted by




where n is three or four;  
a ring denoted by



where  $R_8$  is  
hydrogen, or  
alkyl of from one to four carbon  
atoms; or

5 a ring denoted by



10 where  $R_9$  and  $R_{10}$  are  
hydrogen or alkyl of from one to  
four carbon atoms; and

$R_4$  is  
15 alkyl of from one to four carbon atoms, or  
trifluoromethyl.


7. A process in accordance with Claim 1, wherein

20 X is  $-\text{CH}_2\text{CH}_2-$ ,

$R_1$  is  
phenyl, or  
phenyl substituted by  
25 fluorine,  
chlorine,  
trifluoromethyl,  
alkyl of from one to four carbon  
atoms,  
30 alkoxy of from one to four carbon  
atoms, or  
alkanoyloxy of from two to eight carbon  
atoms;

35  $R_2$  and  $R_3$  are independently  
carboalkoxy of from two to eight carbon  
atoms or,

when taken together with the carbon atoms to which they are attached form a ring denoted by



wherein R<sub>8</sub> is hydrogen or alkyl of from one to four carbon atoms; and R<sub>4</sub> is isopropyl or trifluoromethyl.

10

8. A process according to Claim 1, in which one of the following compounds is prepared : trans-6-[2-[3,4-dichloro-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- 15 trans-6-2-[3,4-dibromo-2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- 20 trans-6-[2-[2-(4-fluorophenyl)-5-(trifluoromethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- trans-dimethyl 2-(4-fluorophenyl)-5-(1-methylethyl)-1-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3,4-dicarboxylate;
- 25 trans-6-[2-[2-(4-fluorophenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- 30 trans-6-[2-[2-cyclopropyl-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;
- 35 trans-6-[2-[2-(1,1-dimethylethyl)-5-(4-fluorophenyl)-1H-pyrrol-1-yl]ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one;

trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxy-phenyl)-5-methyl-1H-pyrrol-1-yl]ethyl]-2H-2-one;  
trans-tetrahydro-4-hydroxy-6-[2-[2-(2-methoxy-phenyl)-5-(1-methylethyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;  
5 trans-tetrahydro-4-hydroxy-6-[2-[2-methyl-5-(1-naphthalenyl)-1H-pyrrol-1-yl]ethyl]-2H-pyran-2-one;  
trans-6-[2-(2-bicyclo[2.2.1]hept-5-en-2-yl-5-methyl-1H-pyrrol-1-yl)ethyl]tetrahydro-4-hydroxy-2H-pyran-2-one; and  
10 trans-6-[2-[2-(4-fluorophenyl)-5-(1-methylphenyl)-1H-pyrrol-1-yl]propyl]tetrahydro-4-hydroxy-2H-pyran-2-one.

15 9. A process for preparing a pharmaceutical composition which process comprises combining a compound prepared in accordance with any preceding claim together with a pharmaceutically acceptable carrier or diluent.

10 10. For use in a method of treatment in which cholesterol biosynthesis in a patient is inhibited, a compound in accordance with any one of Claims 1 to 8 or a pharmaceutical composition in accordance with Claim 9.

25

30

35

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**